

## NAIL AVULSION AFTER CARE

Patient Name: \_\_\_\_\_



### At Home for the first 12-24 hours:

***For Pain: take your pain medication as instructed by the doctor***

IBUPROFEN - Take 600mg (3 x 200mg tablets) every 6 hours  
TYLENOL - Take 1000mg (2 x 500mg tablets) every 6 hours

### 24 Hours After Surgery - The First Soak


Do your first soak on: \_\_\_\_\_

- soak your toe in warm water with 1-2 tablespoons of Epsom Salt in it
- remove the big gauze dressing while your toe is in the water

***After the dressing is off, continue to soak your toe(s) for an additional 15-20 minutes***

#### After Soaking:

- dry your foot/feet and put 2 square pieces of gauze (2x2) on your toe(s)
- secure the gauze snugly by wrapping the roll gauze around the toe 2-3 times
- use tape to hold the roll gauze in place



#### Starting on: \_\_\_\_\_

- the toe(s) must be soaked 2 times EACH DAY for 15-20 minutes EACH TIME until your toe is HEALED (2-4 weeks)

***If you do not soak your toe(s) 2 times each day for 15-20 minutes each time, it will not heal properly (will take longer to heal and look very messy)***

#### FAQs:

- **Can I put cream or ointment on my toe?** *It is not necessary to put any cream/ointment on your toe.*
- **My toe looks a bit red and infected. Is this normal?** *Yes, this is part of the normal healing process - **antibiotics are generally not necessary***
- **When should I contact the doctor who did my surgery if I'm concerned about infection?** *You should contact the doctor if the redness worsens or goes beyond the area around the surgery (i.e. if the redness goes beyond the toe, onto the foot) or if your pain worsens*
- **Can I play sports after my surgery?** *No sports, jogging, etc., until your toe is completely healed (approximately 4 weeks)*
- **Can I shower after the procedure?** *Yes, you can shower after the procedure. Have a shower with the bandages on, then do a 15 minute soak*

**FOLLOW-UP:** \_\_\_\_\_

**If you have a concern with your toe**, contact the doctor who did your surgery.

#### After 1 week:

- stop bandaging the toe(s)
- leave your toe open to the air as much as possible